

Comparison of Models for the Dielectric Constant of Seawater and Fresh Water

The following graphs compare models from various sources for the dielectric constant of water; the emissivity of a flat surface computed from the Fresnel equations is plotted at 275 and 295 K. A linear-with-frequency baseline was subtracted, to allow an expanded vertical scale. The curve labeled FASTEM is a model from S. English which uses the Lamkaouchi et al. dielectric constant, and adds a parameterization for surface roughness. That parameterization is slightly negative at zero wind speed, which is the plotted condition.

The Klein-Swift and Guillou et al. models are single Debye curves and therefore have been limited to < 40 GHz; the other models are double-Debye. Some of the models include dependence on salinity S , and these are compared with the fresh-water model of Liebe et al. in the graphs for $S=0$.

The stand-alone microwave forward model has a modified version of FASTEM that uses the Guillou et al. model (with $S=0.035$) below 50 GHz, and Lamkaouchi et al. at higher frequencies. It does not differentiate between sea water and fresh water.

References

- C. Guillou, W. Ellison, L. Eymard, K Lamkaouchi, C. Prigent, G. Delbos, G. Balana, and S. A. Boukabara, "Impact of new permittivity measurements on sea surface emissivity modeling in microwaves," *Radio Science*, **33**, 649-667 (1998).
- L. A. Klein and C. T. Swift, "An improved model for the dielectric constant of sea water at microwave frequencies," *IEEE Trans Ant. & Prop.* **AP-25**, 104-111 (1977).
- K. Lamkaouchi, G. Balana and W. Ellison, "New Permittivity Data for Sea Water (30-100 GHz)", ESA report 11197/94/NL/CN (1997).
- H. J. Liebe, G. A. Hufford, and T. Manabe, "A model for the complex permittivity of water at frequencies below 1 THz", *Int. J. IR and MM Waves*, **12**, 659-675 (1991).
- A. P. Stogryn, H.T. Bull, K. Rubayi, S. Iravanchy, Aerojet report (1995).







